



FOR INCREASED FOLIAGE AND YIELD

GENERAL INFORMATION

When applied to young growing plants, **INCREASE** directly affects growth by stimulating additional uptake of nutrients and energy. It does so by increasing the foliage produced by the plant. Simply stated, the more foliage available to the sun for photosynthesis, the more productive will be the plant. When used as directed, **INCREASE** should not affect the number of bud sets on the plant.

Increase will increase the thickness and durability of plant foliage insuring that there is plenty of foliage available for photosynthesis to manufacture the complex sugars required to produce quality fruit.

STORAGE

Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in temperatures between 5 C (41 F) and 25 C (77 F).

DISPOSAL

Dispose of Waste Material according to local, state, and Federal environmental laws and regulations.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trail application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water. If irritation persists, obtain medical attention.

GUARANTEED ANALYSIS

MINIMUM GUARANTEED ANALYSIS:

Calcium (Ca)..... 24.0%

Chlorine, not more than ... 46.0%

Derived from: Calcium Chloride humate

Compatibility

Test to ensure compatibility with foreign spray mixes, a jar & Crop to be Sprayed, compatibility test is recommended before mixing with other products other than water in the tank.

APPLICATION

Mix 2 lbs **INCREASE** into sufficient water to cover 10 acres. Spray in a fine mist at high nozzle pressure to ensure coverage of both sides of the leaves. Moisten to the point of run-off. Do not saturate. As a rule, **INCREASE** should be applied when crops are 5 to 10 inches in height (length in vine crops). In fall-planted crops, apply soon after start of re-growth.

Metric:

Apply .25 kgs INCREASE with sufficient water to cover per 1 ha. Spray in a fine mist at highest pressure. Moisten both sides of the leaves only

Always Follow Spray Schedule provided by UAS of America representative.

Manufactured by

UAS of America, Inc.

534 CR 529A Lake Panasoffkee, FL 33538

Tel 1-352-793-1682 Fax 1-352-793-4864

"COMMERCIAL USE ONLY"

Warranty

BUYER ASSUMES ALL RESPONSIBILITY FOR USE, STORAGE, HANDLING AND DISPOSAL OF THIS PRODUCT WHEN NOT IN ACCORDANCE WITH LABEL DIRECTIONS. UAS of AMERICA AND ITS DISTRIBUTOR'S LIMIT OF WARRANTY EXTENDS ONLY TO REPLACEMENT OF DEFECTIVE PRODUCT.